

Smallpox: Frequently Asked Questions

What should I know about smallpox?

Smallpox is an acute, contagious, and sometimes fatal disease caused by the variola virus (an orthopoxvirus), and marked by fever and a distinctive progressive skin rash. In 1980, the disease was declared eradicated following worldwide vaccination programs. However, in the aftermath of the events of September, 2001, the U.S. government is taking precautions to be ready to deal with a bioterrorist attack using smallpox as a weapon. As a result of these efforts: 1) There is a detailed nationwide smallpox preparedness program to protect Americans against smallpox as a biological weapon. This program includes the creation of preparedness teams that are ready to respond to a smallpox attack on the United States. Members of these teams – health care and public health workers – have been vaccinated so that they might safely protect others in the event of a smallpox outbreak. 2) There is enough smallpox vaccine to vaccinate everyone who would need it in the event of an emergency.

What are the symptoms of smallpox?

The symptoms of smallpox begin with high fever, head and body aches, and sometimes vomiting. A rash follows that spreads and progresses to raised bumps and pus-filled blisters that crust, scab, and fall off after about three weeks, leaving a pitted scar.

If someone comes in contact with smallpox, how long does it take to show symptoms? After exposure, it takes between 7 and 17 days for symptoms of smallpox to appear (average incubation time is 12 to 14 days). During this time, the infected person feels fine and is not contagious.

Is smallpox fatal?

The majority of patients with smallpox recover, but death may occur in up to 30% of cases. Many smallpox survivors have permanent scars over large areas of their body, especially their face. Some are left blind.

How is smallpox spread?

Smallpox normally spreads from contact with infected persons. Generally, direct and fairly prolonged face-to-face contact is required to spread smallpox from one person to another. Smallpox also can be spread through direct contact with infected bodily fluids or contaminated objects such as bedding or clothing. Indirect spread is less common. Rarely, smallpox has been spread by virus carried in the air in enclosed settings such as buildings, buses, and trains. Smallpox is not known to be transmitted by insects or animals.

If smallpox is released in aerosol form, how long does the virus survive?

The smallpox virus is fragile. In laboratory experiments, 90% of aerosolized smallpox virus dies within 24 hours; in the presence of ultraviolet (UV) light, this percentage would be even greater.

Is smallpox contagious before the smallpox symptoms show?

A person with smallpox is sometimes contagious with onset of fever (prodrome phase), but the person becomes most contagious with the onset of rash. The infected person is contagious until the last smallpox scab falls off.

Is there any treatment for smallpox?

Smallpox can be prevented through use of the smallpox vaccine. There is no proven treatment for smallpox, but research to evaluate new antiviral agents is ongoing. Early results from laboratory studies suggest that the drug cidofovir may fight against the smallpox virus; currently, studies with animals are being done to better understand the drug's ability to treat smallpox disease. Patients with smallpox can benefit from supportive therapy (e.g., intravenous fluids, medicine to control fever or pain) and antibiotics for any secondary bacterial infections that may occur.

What is the smallpox vaccine, and is it still required?

The smallpox vaccine is the only way to prevent smallpox. The vaccine is made from a virus called *vaccinia*, which is another "pox"-type virus related to smallpox but cannot cause smallpox. The vaccine helps the body develop immunity to smallpox. It was successfully used to eradicate smallpox from the human population.

Routine vaccination of the American public against smallpox stopped in 1972 after the disease was eradicated in the United States. Until recently, the U.S. government provided the smallpox vaccine only to a few hundred scientists and medical professionals who work with smallpox and similar viruses in a research setting. After the events of September, 2001; however, the U.S. government took further actions to improve its level of preparedness against terrorism. For smallpox, this included updating a response plan and ordering enough smallpox vaccine to immunize the American public in the event of a smallpox outbreak. The plans are in place, and there is sufficient vaccine available to immunize everyone who might need it in the event of an emergency. In addition, in December of 2002, the Bush Administration announced a plan to better protect the American people against the threat of smallpox attack by hostile groups or governments. This plan includes the creation of smallpox healthcare teams that would respond to a smallpox emergency. Members of these teams were vaccinated against smallpox. The plan also included vaccination of certain military and civilian personnel who are or may be deployed in high threat areas.

If someone is exposed to smallpox, is it too late to get a vaccination?

Vaccination within 3 days of exposure will completely prevent or significantly modify smallpox in the vast majority of persons. Vaccination 4 to 7 days after exposure likely offers some protection from disease or may modify the severity of disease.

How long does a smallpox vaccination last?

Past experience indicates that the first dose of the vaccine offers protection from smallpox for 3 to 5 years, with decreasing immunity thereafter. If a person is vaccinated again later, immunity lasts longer.